Distal Radius Fracture in Adults Questionnaire

1.	Which hospital do you work at?	
	Glangwili General Hospital, Carmarthen	
2.	Does your hospital accept or manage trauma γ_{ϵ} patients?	es YES No
3.	Is your hospital a designated major trauma centre?	es No NO
4.	How many adults with closed distal radius fracture does your	r hospital manage in a month?
	ESTIMATED 15-20	
5.	Does your department have a written guideline for the invential closed distal radius fracture following trauma?	vestigation and management of Yes No Verbal Only
6.	If yes, where is your guideline taken from? (For example, E etc.)	3OAST, locally derived guideline
	INDIVIDUAL CONSULTATION PRACTICE	
7.	For patients requiring a closed manipulation of their fractuments where is this manipulation carried out?	re that present during the day,
Ī	Plaster room	YES Please specify:
	In the emergency department (excluding resuscitation area)	riease specify.
-	Resuscitation bay in the emergency department	
Ì	Operating Theatre	
Ī	Other (please specify)	
8.	Which specialty is responsible for the <u>initial</u> manipulation of	the fracture?
	Emergency YES Trauma and Orthopaedics Medicine	Other (please specify)
9.	What form of analgesia is most commonly used for the mani	pulation procedure?
	Regional nerve blockade	Please specify:
	Regional intravenous anaesthesia (Bier's block)	, ,
Ĺ	Local haematoma block YES	

Procedural	sedation	and	analgesia	(please	
specify)					
General anaesthetic					
No specific method of analgesia specified					

10. If a Bier's block is performed, which specialty performs the Bier's block?

		 Diameter of the
Emergency Medicine	Anaesthetics	Please specify:
Trauma and Orthopaedics	Other (please specify)	

11. What grade of doctor is most commonly responsible for performing the Bier's Block?

N/A			

12. For patients requiring a closed manipulation of their fracture that present *overnight*, where is this manipulation carried out?

Plaster room	YES		Plea
In the emergency department (excluding resuscitation			
area)			
Resuscitation bay in the emergency department			
Operating Theatre			
Other (please specify)			
Manipulation not carried out overnight		'	

Please specify:	

13. Overnight, which specialty is responsible for the initial manipulation of the fracture?

Emergency Medicine	YES	Manipulation not carried out overnight	
Trauma and Orthopaedics		Other (please specify)	

Please specify:

14. Overnight, what form of analgesia is used for the manipulation procedure?

Regional nerve blockade	
Regional intravenous anaesthesia (Bier's block)	
Local haematoma block	YES
Procedural sedation (please specify)	
General anaesthetic	
No specific method of analgesia specified	
Manipulation not carried out overnight	

Please specify:

15. If patients do not receive manipulation out-of-hours where and when do they return?

Where		When	
Emergency Department		Following morning	
Fracture clinic	YES	Next working day (i.e. on Monday if seen over the weekend)	
Plaster room		Next available routine fracture clinic	

16. Following presentation to hospital with a distal radius fracture, how soon after the injury are patients seen in the Fracture Clinic? VARIES

Within 24 hours	
Within 48 hours	
Within 72 hours	
More than 3 days later	