Paediatric Forearm Fracture Questionnaire

Which hospital do you work at?				
Bronglais General Hospital				
Does your hospital accept or manage paediatric trauma patients?	Yes	Х	No	
Is your hospital a designated major trauma centre?	Yes		No	Х
, , , , , , , , , , , , , , , , , , , ,				
		Bronglais General Hospital Does your hospital accept or manage paediatric trauma patients?	Bronglais General Hospital Does your hospital accept or manage paediatric trauma patients?	Bronglais General Hospital Does your hospital accept or manage paediatric trauma patients? Yes X

- Approximately how many paediatric patients with a closed forearm fracture does your hospital manage in a month?
 Approximately 10 (rough figures; please speak to the coding department if more accurate figures are required).
- 5. Does your department have a written guideline for the investigation and management of potential closed forearm fracture in children?


Yes	
No	
Verbal Only	Х

- 6. If yes, where is your guideline taken from? (For example, NICE, locally derived guideline etc.)
- 7. Does your guideline specify criteria for performing closed reduction in the emergency department of closed forearm fractures? If so, please specify.

No

8. For patients requiring a closed manipulation of their fracture that present *during the day*, <u>where</u> is this manipulation carried out?

Plaster room	
In the emergency department (excluding resuscitation area)	
Resuscitation bay in the emergency department	
Operating Theatre	Х
Other (please specify)	



9. Which specialty is responsible for the initial manipulation of the fracture?

Emergency Medicine	Trauma and Orthopaedics	Х	Other (please specify)	
			Please specify:	

10. What form of analgesia is most commonly used for the manipulation procedure?

Nasal diamorphine and Entonox		Please
Nasal fentanyl and Entonox		
Procedural sedation (please specify)		
General anaesthetic	Х	
No specific method of analgesia specified		
Other (please specify)		

specify:

11. For buckle fractures, what immobilisation, if any, is provided?

Splint	Х
Plaster of Paris	Х
Wool and crêpe bandage	
Other (please specify)	

Please specify:		

12. How are closed, overriding fractures of the distal radius metaphysis managed? (Please answer for both age ranges)

(ricuse unswer jor both uge runges)			
Under 10 years old		10 years old and over	
Moulded cast		Moulded cast	
Formal manipulation (not in theatre)		Formal manipulation (not in theatre)	
Formal manipulation and k-wire fixation		Formal manipulation and k-wire fixation	
(in theatre)		(in theatre)	
Other (please specify)	Х	Other (please specify)	Х
Formal manipulation +/- k-wire fixation (in		Formal manipulation +/- k-wire fixation	(in
theatre)		theatre)	

13. If a patient requires manipulation or treatment in theatre (closed injury, without neurovascular compromise), what is the usual pathway?

Admitted to inpatient ward for next available daytime trauma list	Х	Please specify:
Admitted to inpatient ward for surgery on an emergent basis		
(for example, manipulation performed overnight)		
Patient discharged to attend outpatient clinic prior to definitive treatment		
Patient discharged and added to rolling trauma list		
(no follow-up in clinic prior to treatment)		
Other (please specify)		

14. Does any of your guideline differ out-of-hours (outside 8:00-17:00, Monday – Friday)? If so, please specify:

No

15. Is there a mechanism in your hospital (e.g. audit process) whereby cases requiring revision manipulations/procedures are collected and reviewed?

Yes No X