



GIG
CYMRU
NHS
WALES

Bwrdd Iechyd Prifysgol
Hywel Dda
University Health Board

TRANSMISSION BASED PRECAUTIONS – CONTACT/AIRBORNE/DROPLET PRECAUTIONS POLICY

Policy Number:	353	Supersedes:		Classification	Clinical
Version No	Date of EqIA:	Approved by:	Date of Approval:	Date made Active:	Review Date:
V2	March 2018	Clinical Written Control Documentation Group	04/04/2018	05/04/2018	04/04/2021
		Extended - Infection Prevention Steering Group Chair (IPSSG)	30/03/2021	30/03/2021	30/09/2021
		Infection Prevention Group Chair	20/01/2022	21/01/2022	20/04/2022

Brief Summary of Document:	This Policy describes when patients require isolation from contact, airborne and droplet infections and the standard infection control precautions required to treat them.
Scope:	This Policy should be used by the Infection Prevention team and all healthcare professionals including nurses and doctors. The policy is provided as a reference source for social care managers and social care
To be read in conjunction with:	149 - Hand Hygiene Policy, 151 - Personal Protective Equipment Policy 154 - Management of Linen Policy 187 - Exposure Management including Sharps Injuries 236 – Outbreak Management Policy 258 – Waste Management Policy 354 - Policy Standard Infection Control Precautions (SICPs), , 230 – Policy for the Management of Blood and Body Fluids, 152 – Policy for the Management of Viral Haemorrhagic Fever (VHF). 232 – Control of the Environment/Environmental cleanliness Policy and Procedure.

HYWEL DDA UNIVERSITY HEALTH BOARD

Owning Committee	Infection Control Committee
------------------	-----------------------------

Executive Director:	Mandy Rayani	Job Title	Director of Nursing Quality & Patient Experience
---------------------	--------------	-----------	--

Reviews and updates		
Version no:	Summary of Amendments:	Date Approved:
1	New Policy	4.3.2014
2	3 yearly review	4.4.2018

Glossary of terms

Term	Definition

Keywords	Airborne, contact, droplet, infectious diseases, gloves, aprons, hand hygiene, linen, waste, isolation
----------	--

HYWEL DDA UNIVERSITY HEALTH BOARD

CONTENTS

1. EXECUTIVE SUMMARY/KEY POINTS.....	4
2. INTRODUCTION.....	4
3. POLICY STATEMENT.....	4
4. SCOPE	5
5. AIM	5
6. OBJECTIVES	5
7. WHAT ARE TRANSMISSION BASED PRECAUTIONS?.....	5
7.1. Why are transmission based precautions necessary?	5
7.2. When should Transmission Based Precautions be applied?	5
7.3. How long should Transmission Based Precautions remain in Place?	5
8. CONTACT PRECAUTIONS PROCEDURE.....	6
8.1. What are contact precautions?.....	6
8.2. Standard Infection prevention and control CONTACT precautions	7
9. AIRBORNE PRECAUTIONS PROCEDURE	9
9.1. What are airborne precautions?	9
9.2. Standard Infection prevention and control AIRBORNE precautions.....	10
10. DROPLET PRECAUTIONS PROCEDURE	13
10.1. What are droplet precautions?	13
10.2. Standard Infection prevention and control DROPLET precautions	14
11. CATEGORY 4 INFECTIOUS DISEASES	17
12. ROLES / RESPONSIBILITIES / FUNCTIONS	18
12.1. Chief Executive	18
12.2. Chief Operating Officer (COO) and Director of Nursing, Quality and Patient Experience.....	19
12.3. Executive Director and Senior Managers.....	19
12.4. Assistant Director of Nursing - Infection Prevention and Control	19
12.5. Locality.....	19
12.6. Responsibility of admitting clinician.....	19
12.7. Ward /Senior Nurse / Directorate Nurses.....	19
12.8. All Clinical staff.....	19
13. TRAINING	19
14. IMPLEMENTATION	19
15. FURTHER INFORMATION	20
16. LIST OF POLICIES TO BE READ IN CONJUNCTION WITH THIS POLICY.....	20
17. REVIEW	20
18. APPENDIX 1 – INFECTIOUS AGENTS OR DISEASE WARRANTING TRANSMISSION BASED PRECAUTIONS IN ADDITION TO STANDARD INFECTION CONTROL PRECAUTIONS	21
19. APPENDIX 2 – ORGANISMS AND DISEASES OR HCAI IMPORTANCE AND APPROPRIATE TBPS.....	22
20. APPENDIX 3 – SUMMARY OF PRECAUTIONS TO MINIMISE THE SPREAD OF INFECTIONS BY THE DROPLET ROUTE.....	34
21. APPENDIX 4 - SUMMARY OF PRECAUTIONS TO MINIMISE THE SPREAD OF INFECTIONS BY CONTACT	35
22. APPENDIX 5 - SUMMARY OF PRECAUTIONS TO MINIMISE THE SPREAD OF INFECTIONS BY THE AIRBORNE ROUTE	36

HYWEL DDA UNIVERSITY HEALTH BOARD

1. EXECUTIVE SUMMARY/KEY POINTS

Transmission Based Precautions (TBPs) are categorised according to the route of transmission of the infectious agent such as contact, airborne and droplet.

TBP's are necessary because transmission of specific agents will not be prevented by Standard Infection Control Precautions (SICPs) alone

TBPs are required in all healthcare settings when a patient is known or suspected to be infected/colonised with an infectious agent or an epidemiologically important organism that can spread by the contact, airborne and droplet route.

The duration of TBPs for specific infectious agents spread by contact, airborne or droplet is listed in Appendix 1 (however, this list is not exhaustive)

Contact precautions are a set of infection control measures which are designed specifically to prevent and control the transmission of infectious agents spread by direct or indirect contact to patients and health care workers during provision of care.

Airborne precautions are a set of infection control measures which are designed specifically to prevent and control the transmission of infectious agents spread by small particles in the respirable size range to patients and healthcare workers during provision of care.

Airborne precautions are used to prevent and control infections spread without necessarily having close patient contact via aerosols (<5µm) from the respiratory tract of one individual directly onto a mucosal surface or conjunctivae of another individual. Aerosols penetrate the respiratory system to the alveolar level.

Droplet precautions are a set of infection control measures which are designed specifically to prevent and control the transmission of infectious agents spread by small droplets to patients and healthcare workers during provision of care.

Droplet transmission is defined as the transfer of a large droplet (>5µm) from the respiratory tract of an infected individual directly onto a mucosal surface of conjunctivae of another individual. Due to the comparative large size of the particles it is accepted that droplets when dispelled only travel short distances through the air, e.g. less than 3 feet (1 metre away). The activity, which resulted in the droplet expulsion from the respiratory tract, affects this distance of spread and therefore has to be considered when precautions are being taken.

2. INTRODUCTION

TBPs, in addition to Standard Infection Control Precautions (SICPs), are a set of measures that must be implemented when patients are either suspected or known to be infected with a specific infectious agent, when aiming to prevent and control spread. The purpose of this policy is to ensure that the correct procedures are followed for patients who have a contact, droplet, or airborne infection.

3. POLICY STATEMENT

The commitment of the Health Board is to promote a culture of zero tolerance to any health care associated infection (HCAI), with the ultimate aim of preventing all avoidable infections through use of TBPs.

4. SCOPE

This Policy should be used by the Infection Prevention team and all healthcare professionals including nurses and doctors. The policy is provided as a reference source for social care managers and social care.

5. AIM

It is the intent that this Policy will provide a common, consistent approach to infection prevention and control, and wherever possible prevent transmission of infection within the healthcare setting.

6. OBJECTIVES

For all health care workers to adhere to contact, airborne and droplet transmission based precautions to aid prevention of transmission of infection.

7. WHAT ARE TRANSMISSION BASED PRECAUTIONS?

TBPs are categorised according to the route of transmission of the infectious agent such as contact, airborne and droplet.

7.1. Why are transmission based precautions necessary?

TBPs are necessary because transmission of specific agents will not be prevented by SICPs alone e.g. *Mycobacterium tuberculosis*. SICPs are the minimum set of measures to be applied at all times within a healthcare setting. SICPs should be applied for both recognised and unrecognised sources of infection and are intended to protect the patient, healthcare worker and any visitors/carers.

SICPs must underpin all healthcare activities and should be applied at all times when exposure to blood, other body fluids secretions or excretions (except sweat) non intact skin or mucous membranes may occur. SICPs also apply to equipment or items in the patient's environment that may have become contaminated.

7.2. When should Transmission Based Precautions be applied?

TBPs are required in all healthcare settings when a patient is known or suspected to be infected/colonised with an infectious agent or an epidemiologically important organism that can spread by the contact, airborne and droplet route.

This includes precautions to be taken with those;

- With active infections
- Who are incubating infectious disease
- Who are asymptomatic but suspected to be infectious
- Who are colonised with pathogenic micro-organisms

7.3. How long should Transmission Based Precautions remain in Place?

The duration of TBPs for specific infectious agents spread by contact, airborne or droplet is listed in Appendix 1 (however, this list is not exhaustive). TBPs may need to be lengthened, e.g. for a patient who is immunocompromised due to risk of prolonged shedding of organisms. This decision should be based on the patient's situation, symptoms, and treatment and be guided by the physician and the IPCT.

8. CONTACT PRECAUTIONS PROCEDURE

8.1. What are contact precautions?

Contact precautions are a set of infection prevention and control measures which are designed specifically to prevent and control the transmission of infectious agents spread by direct or indirect contact to patients and healthcare workers during provision of care.

These precautions include;

- Isolation
- Hand hygiene
- Use of personal protective equipment (PPE)
- Care of equipment and environment including decontamination
- Safe handling of linen
- Safe handling of waste

8.1.1. Why are contact precautions important within the health and social care settings?

It is important to prevent infectious agents that could be present on, for example, a patient's skin/mucous membranes, or immediate environment, from being transmitted via contact to others and resulting in HCAs. This is the most common mode of transmission of infectious agents in healthcare settings and therefore it is essential that all healthcare workers understand how to prevent spread via this route

8.1.2. What is the rationale for contact precautions?

Contact precautions are required to prevent the transmission of infectious agents via direct and indirect contact and to minimise HCAs. Due to the nature of contact transmission the precautions described are pivotal and must be applied during any health and social care activity, even those not formally associated with additional infection control precautions. This includes activities such as feeding, playing, or close contact care when a patient has an infectious agent.

- Direct contact transmission is when an infectious agent is transferred directly from one person to another e.g. skin to skin contact when scabies is present or transfer of an infectious agent from an open wound of an infected individual to the mucous membranes or skin break in another susceptible individual.
- Indirect contact transmission is when an infectious agent is transferred to an individual from an object and/or other person. This can occur in a number of ways and varies depending on the infectious agent. An example of a mode on indirect transmission includes via the hands of healthcare workers after contact with an infected or colonized patient's environment, patient care equipment, or surgical instruments which have been inadequately decontaminated. In addition, transmission can occur through direct contact via toys.

8.1.3. When are contact precautions required?

The need for contact precautions will vary depending on the patient, the setting and the infectious agent (please refer to Appendix 1) and the procedures/activities being undertaken. For example, highly dependent in-patient areas may require different considerations to a mental health setting.

Contact precautions are required in all health care settings when a patient is known or suspected to be infected/colonized with certain infectious agents or epidemiologically important organism that can be spread by contact. These precautions are to be taken when patients have active infections, those who are incubating an infectious disease and those who are asymptomatic but suspected to be infectious and those who are colonized with pathogenic micro-organisms.

HYWEL DDA UNIVERSITY HEALTH BOARD

8.2. **Standard Infection prevention and control CONTACT precautions**

8.2.1. Patient placement/isolation

Patient placement or isolation requires a risk assessment in order to determine the most appropriate placement for the patient. This will depend on;

- The infectious agent
- The patient and their overall condition e.g. a productive cough
- The area the patient is being cared for. This includes the potential for adverse outcomes in others and the availability of single rooms.
- The procedure/activities being undertaken

Where appropriate, single rooms with ensuite toilet and hand washing facilities are preferred for patients with known/suspected infections requiring contact precautions. The requirement to keep the door shut shall be on a risk assessment basis but is considered good practice.

If placement in a single room with toilet and hand washing facilities or transfer is not possible, placement decisions should be subject to a risk assessment and discussion with the IPCT. Placement might include;

- In a single room with no ensuite facilities
- Cohorting – where cohorting is the only option this should be based on placing those with the same known/suspected infection in a designated area. If cohorting is not possible, do not place with immunocompromised patients
- The decision to either cohort or use rooms/cubicles without facilities will depend on a number of factors. Advice should be sought from the IPCT.

8.2.2. Hand Hygiene

Hand hygiene is essential in preventing HCAs particularly for contact precautions. Please refer to the Health Board 149 - Hand Hygiene Policy.

8.2.3. Respiratory Hygiene and Cough Etiquette

Reinforce good respiratory hygiene with the patient at all times and assist as necessary

- Ensure patients cover the mouth and nose with a disposable tissue.
- Wearing gloves, place tissue into a waste bin.
- Remove gloves and wash hands.

8.2.4. Personal Protective Equipment PPE

PPE is essential in preventing HCAs particularly for contact precautions. Please refer to the Health Board 151 - Personal Protective Equipment Policy.

Gloves and aprons;

- Disposable gloves and plastic aprons should be put on and worn during care activities and where there will be contact with the patient or their immediate environment.
- The use of a disposable fluid repellent gown maybe more appropriate in order to gain more protection including the arms for specific infections. Advice should be sought from the IPCT.

Face Protection;

- Face/eye protection including masks and goggles are required if there is a risk of mucosal splashing to the eyes and mouth.

Good practice points;

- Supplies of PPE must be available at the entrance to single or cohort rooms.
- Aprons and gloves must be put on before undertaking care activities.

HYWEL DDA UNIVERSITY HEALTH BOARD

- No outer coats to be worn.
- PPE must be removed immediately upon leaving the room followed by hand washing.
- PPE must be changed between different procedures and care activities including gloves.
- PPE must be changed and hand hygiene performed between contact with every patient/client/resident, including others being cared for under contact precautions within the same area.
- Safe disposal of PPE is essential immediately following removal.

8.2.5. Management of Care Equipment

Care equipment must be given additional consideration in order to prevent the spread of infectious agents that might be contaminating items.

- Equipment must be allocated to individuals being cared for under contact precautions e.g. commodes.
- Equipment must not be shared with others unless thoroughly decontaminated first.
- Items of equipment must be intact. Items that are not intact must be removed and replaced with intact items.
- Where possible use single use only disposable products.

8.2.6. Control of the Environment

Care of the environment must be given additional consideration in order to prevent the spread of infectious agents that might be contaminating items. Please refer to the Health Board 232 – Control of the Environment/Environmental cleanliness Policy and Procedure.

- The environment must be cleaned at least daily and when visibly contaminated. Particular attention must be given to frequently touched items e.g. door handles, bed tables etc.
- An increase in frequency must be considered particularly during outbreaks of infection e.g. diarrhoea and/or vomiting
- The environment must be clutter free to allow for effective cleaning
- Equipment for cleaning must follow the Health Board's colour coded cleaning system. These items must be clean, fit for purpose and decontaminated and/or disposed of appropriately
- Terminal cleaning of the environment MUST be performed prior to use by any other patient.

8.2.7. Safe Management of Linen

Linen that could be contaminated must be managed safely in order to avoid cross transmission of infectious agents. They should be bagged following the Health Board 154 - Safe Management of Linen Policy as follows;

- Place contaminate linen into an alginate bag at the point of removal. DO NOT carry linen out of the room.
- Place alginate bag in appropriate colour coded bag.
- Remove gloves and wash hands.
- Communicate with others who may handle linen to ensure they take appropriate precautions.

8.2.8. Management of blood and body fluid spillage

All body fluid spillages must be cleaned and decontaminated following the Health Board 230 - Management of Body Fluid Policy.

HYWEL DDA UNIVERSITY HEALTH BOARD

8.2.9. Safe Management of Waste

All waste must be segregated and disposed of in accordance with 258 – Waste Management Policy.

- Waste that could be contaminated must be managed safely in order to avoid cross infection of infectious agents.
- All waste generated from an infected or suspected of being infected patient must be disposed of into orange clinical waste bags (acute setting) or appropriate colour waste bags for contaminated waste.

9. **AIRBORNE PRECAUTIONS PROCEDURE**

9.1. **What are airborne precautions?**

Airborne precautions are a set of infection control measures which are designed specifically to prevent and control the transmission of infectious agents spread by small particles in the respirable size range to patients and healthcare workers during provision of care.

These precautions include;

- Patient placement/Isolation
- Respiratory hygiene/cough etiquette
- Hand hygiene
- Use of personal protective equipment (PPE)
- Care of equipment and environment including decontamination
- Management of body fluid spillage
- Cleanliness of the environment
- Safe handling of linen
- Safe handling of waste
- Occupational exposure management (including sharps injuries).

9.1.1. Why are airborne precautions important within the health care settings?

It is important to prevent infectious agents that could be present in, for example, the respiratory tract of individuals, being transmitted via contact to others and resulting in HCAs.

9.1.2. What is the rationale for airborne precautions?

Airborne precautions are used to prevent and control infections spread without necessarily having close patient contact via aerosols (<5µm) from the respiratory tract of one individual directly onto a mucosal surface or conjunctivae of another individual. Aerosols penetrate the respiratory system to the alveolar level.

It is essential to apply the relevant infection control precautions during any relevant health care activity. Due to the fact that these are small particles that remain infectious over time and distance and are able to enter the respiratory tract of others without necessarily having close contact (or being in the same room), the precautions described are pivotal. Any aerosol generating procedure which results in the expulsion of small particles within the respiratory size range must be considered when precautions are taken.

HYWEL DDA UNIVERSITY HEALTH BOARD

Aerosol generating procedures include: -

- Intubation/Extubation
- Nasopharyngeal aspiration
- Tracheostomy care
- Chest physiotherapy
- Bronchoscopy
- Continuous positive airways pressure (PAP)
- Non-invasive ventilation
- Suctioning
- Humidification.

FFP3 Masks or equivalent to be worn (Health Protection Agency 2012)

Infected individuals can cause respiratory droplets to be expelled as a result of coughing, sneezing and even talking. There can also be a risk during certain healthcare procedures such as endotracheal intubation and suctioning.

9.1.3. When are airborne precautions required?

The need for airborne precautions will vary depending on the patient, the setting and the infectious agent (please refer to Appendix 1) and the procedures/activities being undertaken. For example, highly dependent in-patient areas may require different considerations to a mental health setting.

Airborne precautions are required in all health care settings when a patient is known or suspected to be infected/colonized with certain infectious agents or epidemiologically important organism that can be spread by the airborne route.

These include precautions to be taken with those with active infections, who are incubating an infectious disease, who are asymptomatic but suspected to be infectious and those who are colonized with pathogenic micro-organisms

9.2. Standard Infection prevention and control AIRBORNE precautions

9.2.1. Patient placement/isolation

Patient placement or isolation requires a risk assessment in order to determine the most appropriate placement for the patient/clone/resident. This will depend on;

- The infectious agent.
- The patient and their overall condition e.g. a productive cough.
- The area the patient is being cared for. This includes the potential for adverse outcomes in others and the availability of single rooms.
- The procedure/activities being undertaken.

Patients with certain known/suspected infections requiring airborne precautions should be placed in an isolation room or negative pressure isolation room with hand washing, toilet and shower (ensuite) facilities as soon as possible. The door to this room MUST remain closed.

If there are no isolation rooms available then placement decisions should be performed with local risk assessments with support from the IPCT and Consultant Microbiologist and may include;

- Placement in a room with ensuite facilities, Door MUST be closed.
- Placement in a single room with no ensuite facilities. The door MUST remain closed. The room must be ventilated if possible via an open window to allow for the exchange of air. The patient should wear a surgical mask when in close contacts with others.
- Cohorting is not recommended for those infections spread via the airborne route.

9.2.2. Hand Hygiene

Hand hygiene is essential particularly for airborne precautions. Please refer to the Health Board 149 – Hand Hygiene Policy.

9.2.3. Respiratory Hygiene and Cough Etiquette

Reinforce good respiratory hygiene with the patient at all times and assist as necessary;

- Cover nose and mouth with disposable single-use tissues when sneezing, coughing, wiping or blowing noses.
- Dispose of all tissues immediately into a orange clinical waste bin.
- Offer and encourage the patient to wash hands after coughing, sneezing, using tissues, or after contact with respiratory secretions and contaminated objects.
- Keep hands away from the mucous membranes of the eyes and nose. Certain patients (e.g. the elderly and children) may need assistance with containment of respiratory secretion; those who are immobile will need a receptacle (e.g. a small plastic orange clinical waste bag) readily at hand for the immediate disposal of used tissues and should be offered hand hygiene facilities.
- Instruct patient and any visitors or carers on the steps described above.

9.2.4. Personal Protective Equipment (PPE)

PPE is essential in preventing HCAs particularly for contact precautions. Please refer to the Health Board 151 - Personal Protective Equipment Policy.

Masks and other facial protection:

- Respiratory masks i.e. FFP3 masks (not surgical masks) are designed to prevent inhalation of infectious airborne particles and subsequent access to the mucous membranes of the respiratory tract of an individual. This is one of the key precautions to be considered when delivering care to those infections transmissible by the airborne route.
- The use of these masks applies to those with active respiratory multi-drug resistant (MDR) *Mycobacterium Tuberculosis*. It also applies when extra pulmonary TB and/or infectious TB lesions are present and aerosol-generating procedures are being undertaken (for advice when FFP3 are no longer required please contact IPCT)

The FFP3 masks must be:

- Fit tested – the efficiency of the mask depends on them being a tight fit to the wearer's face. This procedure must be performed by either trained Health & Safety representative or IPCT.
- Fit checked – each time a FFP3 mask has been put on it should be fit checked by the wearer before entering the patient's area.
- Put on before entering the affected area.
- Changed if you feel you cannot breathe with it on or it is damaged or torn by you (you should leave the room immediately and only then removed and dispose of the mask).
- not be routinely worn when caring for those patients with chickenpox, measles or disseminated herpes zoster. This is due to the fact that once the skin lesions are evident and the infectious agent is known to be present, communicability through the airborne route is less significant. This is known to be the case because it is expected that most health care workers have immunity through vaccination/exposure to chickenpox/measles.
- Worn by patients due to the nature of the mask which filters inhaled and not exhaled air.

HYWEL DDA UNIVERSITY HEALTH BOARD

Surgical Masks:

The use of surgical masks in other situations where respiratory secretions might be spread by the airborne route must be considered along with the use of face protection following SICPs. Surgical masks being worn must be changed when contaminated or wet with breath moisture or if damaged/torn. Other face protection, e.g. eye protection provided by goggles or visors, may be required if there is a risk of mucosal splashing to the face as a result for example, coughing, sneezing or aerosol generating procedures.

Gloves and aprons:

- Disposable gloves and plastic aprons must be put on and worn during care activities and where there will be contact with the patient or their immediate environment
- The use of a disposable fluid repellent gown maybe more appropriate in order to gain more protection including the arms for specific infections. Advice should be sought from the IPCT.

•

Good practice points:

- Supplies of PPE must be available at the entrance to single or cohort rooms.
- Aprons and gloves must be put on before undertaking care activities.
- No outer coats to be worn.
- PPE must be removed immediately upon leaving the room followed by hand washing.
- PPE must be changed between different procedures and care activities including gloves.
- PPE must be changed and hand hygiene performed between contacts with every patient, including other patientd being cared for under contact precautions within the same area.
- Safe disposal of PPE is essential immediately following removal.

9.2.5. Management of Care Equipment

Care equipment must be given additional consideration in order to prevent the spread of infectious agents that maybe contaminating items.

- Equipment must be allocated to individuals being cares for under airborne precautions e.g. blood pressure cuffs.
- Equipment must not be shared with others unless effectively decontaminated first.
- Items of equipment must be intact. Items that are not intact must be removed and replaced with intact items.
- Where possible use single use only disposable products.

9.2.6. Control of the Environment

Care of the environment must be given additional consideration in order to prevent the spread of infectious agents that might be contaminating items. Please refer to the Health Board 232 – Control of the Environment/Environmental cleanliness Policy and Procedure.

- The environment must be cleaned at least daily and when visibly contaminated. Particular attention must be given to frequently touched items e.g. door handles, bed tables etc.
- An increase in frequency must be considered particularly if patients are producing copious amounts of respiratory secretions.
- The environment must be clutter free to allow for effective cleaning.
- Equipment for cleaning must follow the Health Board colour coded cleaning system. These items must be clean, fit for purpose and decontaminated and/or disposed of appropriately.

HYWEL DDA UNIVERSITY HEALTH BOARD

- Terminal cleaning of the environment **MUST** be performed prior to use by any other patient.

9.2.7. Safe Management of Linen

Linen that could be contaminated must be managed safely in order to avoid cross transmission of infectious agents. They should be bagged following the Health Board 154 - Management of Linen Policy;

- Place contaminate linen into an alginate bag at the point of removal. **DO NOT** carry linen out of the room.
- Place alginate bag in appropriate colour coded bag.
- Remove gloves and wash hands.
- Communicate with others who may handle linen to ensure they take appropriate precautions.

9.2.8. Management of blood and body fluid spillage

All body fluid spillages must be cleaned and decontaminated following the Health Board 230 - Management of Body Fluid Policy.

9.2.9. Safe Management of Waste

All waste must be segregated and disposed of in accordance with 258 – Waste Management Policy.

- Waste that could be contaminated with small particles in the respirable size range must be managed safely in order to avoid cross infection.
- All waste generated from an infected or suspected of being infected patient must be disposed of into orange clinical waste bags (acute setting) or appropriate colour waste bags for contaminated waste.

9.2.10 Occupational exposure management

Occupational exposure management is essential for the protection of healthcare workers and relevant immunisation is an essential consideration when caring for those individuals with diseases spread by the airborne route;

- Ensure occupational immunizations are up to date
- Report and manage occupational exposure incidents immediately
- Non immune healthcare workers and those who are pregnant should not provide any close care for individual with specific infectious agents transmitted via the airborne route e.g. chickenpox and measles. Advice must be sought from Occupational Health for immunization, exclusion advice and post exposure guidance

10. **DROPLET PRECAUTIONS PROCEDURE**

10.1. **What are droplet precautions?**

Droplet precautions are a set of infection control measures which are designed specifically to prevent and control the transmission of infectious agents spread by small droplets to patients and healthcare workers during provision of care.

These precautions include;

- Patient placement/Isolation
- Respiratory hygiene/cough etiquette
- Hand hygiene
- Use of personal protective equipment (PPE)
- Care of equipment and environment including decontamination
- Management of body fluid spillage

HYWEL DDA UNIVERSITY HEALTH BOARD

- Cleanliness of the environment
- Safe handling of linen
- Safe handling of waste
- Occupational exposure management (including sharps injuries).

10.1.1. Why are droplet precautions important within the health and social care settings?

It is important to prevent infectious agents that could be present in, for example, the respiratory tract of individuals being transported via droplets to others and resulting in HCAs.

10.1.2. What is the rationale for droplet precautions?

Droplet precautions are required to prevent the transmission of infectious agents via droplets and to minimize HCAs. It is essential to apply the relevant infection control precautions during any relevant health and social activity. Due to the distance that droplets can travel from infected respiratory tracts, which depends on a number of factors including their size, speed, density and a number of additional environmental factors such as temperature, humidity etc, the precautions described are pivotal. Droplet transmission is defined as the transfer of a large droplet (>5um) from the respiratory tract of an infected individual directly onto a mucosal surface or conjunctivae of another individual. Due to the comparative large size of the particles it is accepted that droplets when dispelled only travel short distances through the air, e.g. less than 3 feet (1 metre away). This distance has been used as an effective measure and prevention measures based on this have been shown to be effective. The activity, which resulted in the droplet expulsion from the respiratory tract, affects this distance of spread and therefore has to be considered when precautions are being taken.

Infected individuals can cause respiratory droplets to be expelled as a result of a number of human activities such as coughing, sneezing and even talking. They can also be a risk during certain healthcare procedures such as endotracheal intubation and suctioning.

10.1.3. When are droplet precautions required?

The need for droplet precautions will vary depending on the patient/client/residents, the setting and the infectious agent (please refer to Appendix 1) and the procedures/activities being undertaken.

For example, highly dependent in-patient areas may require different considerations to a mental health setting.

Droplet precautions are required in all health and social care settings when a patient/client/resident is known or suspected to be infected/colonized with certain infectious agents or epidemiologically important organism that can be spread by the droplet route.

These include precautions to be taken with those with active infections, who are incubating an infectious disease, who are asymptomatic but suspected to be infectious and those who are colonized with pathogenic micro-organisms

10.2. **Standard Infection prevention and control DROPLET precautions**

10.2.1. Patient placement/isolation

Patient placement/isolation requires a risk assessment in order to determine the most appropriate placement for the patient. This will depend on;

- The infectious agent
- The patient and their overall condition e.g. a productive cough

HYWEL DDA UNIVERSITY HEALTH BOARD

- The area the patient is being cared for. This includes the potential for adverse outcomes in others and the availability of single rooms.
- The procedure/activities being undertaken.

Patients with certain known/suspected infections requiring droplet precautions should be placed in a single room with ensuite facilities as soon as possible. The door to this room MUST remain closed.

NB Certain conditions may require specialist monitoring and specialist isolation rooms e.g. Severe Acute Respiratory Syndrome (SARS).

If there are no specialised isolation rooms available then placement decisions must be performed with local risk assessments with support from the IPCT and Consultant Microbiologist and may include;

- Placement in a room with ensuite facilities. Door MUST be closed
- Placement in a single room with no ensuite facilities. The door MUST remain closed.
- The room must be ventilated if possible via an open window to allow for the exchange of air. The patient should wear a surgical mask when in close contacts with others.
- Cohorting – where cohorting is the only option this should be considered based on placing those with the same known/suspected infection in the same designated area. This approach is particularly relevant when there are increased numbers of cases e.g. influenza
- If cohorting those with the same infection is not possible avoid placing the infected patient with those who are immunocompromised or within long stay facilities such as care homes, IPCT will advise
- Cohorted patients must be at least 3 feet away from each other.

10.2.2. Hand Hygiene

Hand hygiene is essential particularly for droplet precautions. Please refer to the Health Board 149 - Hand Hygiene Policy.

10.2.3. Respiratory Hygiene and Cough Etiquette

Reinforce good respiratory hygiene with the patient at all times and assist as necessary;

- Cover nose and mouth with disposable single-use tissues when sneezing, coughing, wiping or blowing noses.
- Dispose of all tissues immediately into a orange clinical waste bin.
- Offer and encourage patient to wash hands after coughing, sneezing, using tissues, or after contact with respiratory secretions and contaminated objects.
- Keep hands away from the mucous membranes of the eyes and nose. Certain patients(e.g. the elderly and children) may need assistance with containment of respiratory secretion; those who are immobile will need a receptacle (e.g. a small plastic orange clinical waste bag) readily at hand for the immediate disposable of used tissues and should be offered hand hygiene facilities
- Instruct all affected patients and any visitors or carers on the steps described.

10.2.4. Personal Protective Equipment PPE

PPE is essential in preventing HCAs particularly for contact precautions. Please refer to the Health Board 151 - Personal Protective Equipment Policy.

Masks and other facial protection:

HYWEL DDA UNIVERSITY HEALTH BOARD

- A surgical mask is one of the key precautions to be considered when providing care in close contact. It should be put on before care is provided i.e. on entry into the room or cohort area.
- Masks are not expected to be routinely worn, e.g. when the health care worker has known or proven immunity, or close contact care is not being provided.
- Respiratory masks (FFP3) are not generally expected to be worn for droplet precautions. These masks would only be required when delivering care to those with additional epidemiological significance and/or transmissible by the airborne route. Please contact IPCT for advice.
- Face/eye protection is required if there is a risk of mucosal splashing to the eyes as a result of coughing/sneezing.

Gloves and aprons:

- Disposable gloves and plastic aprons should be put on and worn during care activities and where there will be contact with the patient/client/resident or their immediate environment.
- The use of a disposable fluid repellent gown might be more appropriate in order to gain more protection including the arms for specific infections. Advice should be sought from the IPCT.

Good practice points:

- Supplies of PPE must be available at the entrance to single or cohort rooms
- Aprons and gloves must be put on before undertaking care activities
- No outer coats to be worn
- PPE must be removed immediately upon leaving the room followed by hand washing
- PPE must be changed between different procedures and care activities. This is particularly important for gloves.
- PPE must be changed and hand hygiene performed between contact with every patient, including others being cared for under contact precautions within the same area.
- Safe disposal of PPE is essential immediately following removal.

10.2.5. Management of Care Equipment

Care equipment must be given additional consideration in order to prevent the spread of infectious droplets/respiratory secretions that may be contaminating items.

- Equipment must be allocated to individuals being cared for under contact precautions e.g. blood pressure cuffs
- Equipment must not be shared with others unless thoroughly decontaminated first
- Items of equipment must be intact. Items that are not intact must be removed and replaced with intact items.
- Where possible use single use only disposable products

10.2.6. Control of the Environment

Care of the environment must be given additional consideration in order to prevent the spread of infectious agents that might be contaminating items. Please refer to the Health Board 232 – Control of the Environment/Environmental cleanliness Policy and Procedure.

- The environment must be cleaned at least daily or when visibly contaminated. Particular attention must be given to frequently touched items e.g. door handles, bed tables etc.
- An increase in frequency must be considered particularly if patients are producing copious amounts of respiratory secretions
- The environment must be clutter free to allow for effective cleaning

HYWEL DDA UNIVERSITY HEALTH BOARD

- Equipment for cleaning must follow the Health Board Colour coded Cleaning system. These items must be clean, fit for purpose and decontaminated and/or disposed of appropriately
- Terminal cleaning of the environment MUST be performed prior for use by any other patients/clients/residents.

10.2.7. Safe Management of Linen

Linen that could be contaminated must be managed safely in order to avoid cross transmission of infectious agents. They should be bagged following the Health Board 154 - Safe Management of Linen Policy.

- Place contaminate linen into an alginate bag at the point of removal. DO NOT carry linen out of the room.
- Place alginate bag in appropriate colour coded bag.
- Remove gloves and wash hands.

Communicate with others who may handle linen to ensure they take appropriate precautions.

10.2.8. Management of blood and body fluid spillage

All body fluid spillages must be cleaned and decontaminated following the Health Board 230 - Management of Body Fluid Policy.

10.2.9. Safe Management of Waste

All waste must be segregated and disposed of in accordance with 258 – Waste Management Policy.

- Waste that could be contaminated must be managed safely in order to avoid cross infection of infectious agents
- All waste generated from an infected or suspected of infected patients should be disposed of into orange clinical waste bags (acute setting) or appropriate colour waste bags for contaminated waste.

10.2.10. Why is occupational exposure management an additional consideration for airborne precautions?

Occupational exposure management is essential for the protection of healthcare workers and relevant immunisation is an essential consideration when caring for those individuals with diseases spread by infectious droplets;

- Ensure occupational immunizations are up to date e.g. influenza
- Report and manage occupational exposure incidents immediately, including any mucosal splashing

11. **CATEGORY 4 INFECTIOUS DISEASES**

The Consultant Microbiologist and the IPCT must be informed immediately if a patient has a suspected or diagnosed with a Category 4 Infectious Disease. The Advisory Committee on Dangerous Pathogens ACDP Hazard Group 4 VHF viruses are;

ARENAVIRADAE	BUNYAVIRIDAE
<u>Old World arenaviruses;</u> <ul style="list-style-type: none">• Lassa• Lujo New World arenaviruses;	<u>Nairoviruses</u> <ul style="list-style-type: none">• Crimean Congo haemorrhagic fever

HYWEL DDA UNIVERSITY HEALTH BOARD

<ul style="list-style-type: none"> • Chapare • Guananto • Junin • Machupo • Sabia 	
<p style="text-align: center;">FLAVIVIRIDAE</p> <ul style="list-style-type: none"> • Kyasanur forest disease • Alkhuma haemorrhagic fever • Omsk haemorrhagic fever 	<p style="text-align: center;">FILOVIRIDAE</p> <ul style="list-style-type: none"> • Ebola • Marburg

The patient must have strict isolation and must be transferred as soon as possible to a High Security Infectious Disease Unit (London or Newcastle) shortly after diagnosis, or on high suspicion of such an infectious disease.

- Royal Free Hampstead NHS Trust London – Telephone (24hrs, ask for Infectious disease Physician on call) 020 7794 0500 Or 0844 8480700 (local rate number for when calling from outside London)
- The Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle – Telephone (24 hrs, ask for Infectious Diseases Physician on call 0191 233 6161

Reference Laboratories for VHF screen

- Microbiology laboratories – Porton Down
Health Protection Agency
Porton Down
Salisbury
Wiltshire
SP4 0JG
Telephone; 01980 612100
- Microbiology Services division – Colindale
62 Colindale Avenue
Colindale
London
NW9 5HT
Telephone 0208 200 4400 or 0208 200 6858 (24 hours).

12. ROLES / RESPONSIBILITIES / FUNCTIONS

It is important that the following key staff understand their individual roles in promoting compliance with TBPs.

12.1. Chief Executive

The Chief Executive has ultimate responsibility for infection prevention and control within Hywel Dda Health Board. This responsibility is delegated to the Director of Nursing and Midwifery.

HYWEL DDA UNIVERSITY HEALTH BOARD

12.2. Chief Operating Officer (COO) and Director of Nursing, Quality and Patient Experience

The Director of Nursing, Quality and Patient Experience has delegated responsibility for Infection Prevention in the Health Board and along with COO must be familiar with this policy and support the implementation of the policy throughout the organisation.

12.3. Executive Director and Senior Managers

The Director of Nursing, Quality and Patient Experience has delegated responsibility for infection prevention and control in the Health Board and along with COO must be familiar with this policy and support the implementation of the policy throughout the organisation.

12.4. Assistant Director of Nursing - Infection Prevention and Control

Operational responsibility for infection prevention and control within the Health Board lies with the Assistant Director Infection Prevention & Control who is responsible for ensuring that this policy is available to staff and processes for monitoring compliance are in place.

12.5. Locality

The Locality IPT will promote implementation of this policy in clinical practice and will conduct regular compliance audits for feedback to wards/departments and Locality management teams.

12.6. Responsibility of admitting clinician

The admitting clinician is responsible for initial communication of the case, in particular to:

-
- Consultant in Communicable Disease (CCDC)

12.7. Ward /Senior Nurse / Directorate Nurses

Ensure all staff are familiar with this policy and ensure the policy is complied with. It is the responsibility of the person in charge to ensure that the care area is safe for practice and this includes environmental cleanliness/maintenance. The person in charge has the authority to act if this is deficient.

12.8. All Clinical staff

All health care workers are required to be familiar with this policy and comply with its contents and are responsible for informing the IPT and their manager immediately of any concerns related to poor compliance.

13. TRAINING

Infection Prevention and Control Training is mandatory every 3 years and contents of this policy are included in this training. Infection Prevention and Control staff perform this training and keep attendance records; however, it is the line managers who are responsible to ensure ALL staff attend this training at the required time.

14. IMPLEMENTATION

Implementation of policies and procedures can only be effective if adequate evaluation and monitoring is used to check the system and ensure any shortcomings are identified and dealt with. Locally, Managers are responsible for initiating an ongoing monitoring process within their areas of responsibility.

HYWEL DDA UNIVERSITY HEALTH BOARD

From an organisation perspective, the Infection Prevention and Control Committee shall be responsible for monitoring that this policy and that appropriate actions are being taken to maintain patient safety.

15. FURTHER INFORMATION

Health Protection Scotland – Transmission Based Precautions Policy 2012

National Institute for Clinical Excellence 2012 'Infection: prevention and control of healthcare-associated infections in primary and community care: Clinical Guideline methods, evidence and recommendations'.

National Clinical Guideline Centre at The Royal College of Physicians; London

16. LIST OF POLICIES TO BE READ IN CONJUNCTION WITH THIS POLICY

The practice recommendations set out are drawn from appraisals of the available professional literature on infection prevention and control, conducted by colleagues at Health Protection Scotland which can be found via the link to the Health Protection Scotland (HPS) web site

<http://www.hps.scot.nhs.uk/HCAlic/ic/standardinfectioncontrolprecautions-sicps.aspx>

This document should be read in conjunction with the following Health Board Policies:

149 - Hand Hygiene Policy,

151 - Personal Protective Equipment Policy

154 - Management of Linen Policy

187 - Exposure Management including Sharps Injuries

236 – Outbreak Management Policy

258 – Waste Management Policy

354 - Policy Standard Infection Control Precautions (SICPs), ,

230 – Policy for the Management of Blood and Body Fluids,

152 – Policy for the Management of Viral Haemorrhagic Fever (VHF).

232 – Control of the Environment/Environmental cleanliness Policy and Procedure.

17. REVIEW

This Policy will be reviewed after 3 years, or sooner, as required.

HYWEL DDA UNIVERSITY HEALTH BOARD

18. APPENDIX 1 – INFECTIOUS AGENTS OR DISEASE WARRANTING TRANSMISSION BASED PRECAUTIONS IN ADDITION TO STANDARD INFECTION CONTROL PRECAUTIONS

Infection Prevention & Control

Transmission Based Precautions (TBPs)

Contact Precautions

- Abscess
- *C.difficile* (consider Airborne route if patient has explosive diarrhoea and excessive flatulence)
- Diarrhoea with suspected infective cause
- Diphtheria (cutaneous)
- Epiglottitis
- Hepatitis type A & E
- Herpes Simplex Virus
- Human metapneumovirus
- Impetigo
- Lice – head, body, pubic
- Multi-drug resistant organism(skin/body fluid)
- Scabies
- Staphylococcal disease – skin, wound, burns
- Herpes Zoster Virus (shingles)

Contact and / or Droplet

- Adenovirus – resp infection in paediatrics
- Bronchiolitis
- Diphtheria – pharyngeal
- Herpes zoster (shingles)
- Influenza – seasonal / pandemic
- Whooping cough
- Meningitis
- Mumps
- Multi-drug resistant organism (respiratory)
- Viral haemorrhagic fevers
- Mycoplasma atypical pneumonia
- Parvovirus B19
- Pneumonia – adenovirus
- Rubella
- Respiratory syncytial virus
- Rotavirus
- Streptococcus group A disease i.e. erysipelas, tonsillitis, scarlet fever, necrotising fascitis, puerperal fever

Airborne

- Measles
- Varicella (chickenpox)
- Norovirus
- Multi-drug resistant organism (respiratory)
- Mycobacterium tuberculosis- pulmonary
- Severe acute respiratory syndrome (SARS)

NB. Consider *C.difficile* in this transmission category if patients have explosive diarrhoea and excessive flatulence.

HYWEL DDA UNIVERSITY HEALTH BOARD

19. APPENDIX 2 – ORGANISMS AND DISEASES OR HCAI IMPORTANCE AND APPROPRIATE TBPS

NB for all Vaccination and Immunisation advice, please refer to the Green Book, Public Health England, 2014 - Christine, can we put in hyperlink please

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
Adenovirus	Respiratory tract infection	Droplet	✓			✓	For AGP* only		Requirements of precautions may be extended due to prolonged shedding* of virus, generally until 48 hours following cessation of symptoms
	Conjunctivitis	Contact							
<i>Bordetella Pertussis</i>	Whooping cough	Droplet	✓			✓ Until patient has completed 5 days of antibiotics	For AGP* only	✓	<p>PHE guidance</p> <ul style="list-style-type: none"> http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/WhoopingCough/ <p>Until 5 days of the commencement of antibiotic therapy. If untreated patient infectious for up to 3 weeks. Post exposure prophylaxis for household contacts and in rare circumstances may be indicated for HCWs following prolonged exposure to respiratory secretions.</p>

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
<i>Chlamydia pneumoniae</i>	Pneumonia	Droplet	✓			✓	For AGP * only		
<i>Clostridium difficile</i>	<i>Clostridium difficile</i> associated disease	Contact	✓						<ul style="list-style-type: none"> PHE C diff management link - http://www.hpa.org.uk/webc/HPAwebFile/HPAwebC/1317138914904 HPA Good Practice – <p>Precautions remain until patient is 48 hours symptom free.</p> <p>Specimens are not required for clearance</p>
Coronavirus	Respiratory tract infection	Droplet	✓			✓	For AGP * only	✓	

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
<i>Corynebacterium diphtheriae/ulcerans</i>	Cutaneous diphtheria	Contact	✓					✓	
	Pharyngeal diphtheria	Droplet	✓			✓	For AGP * only	✓	Until negative. Patient considered negative when two cultures (Nasal Pharyngeal Aspirate) are taken 24 hours apart.
GI infections E.g. Salmonella Campylobacter		Contact	✓					✓	Varies depending on the organism seek local guidance from local Infection Control/Health Protection team. Many enteric pathogens are notifiable diseases consult local Health Protection Team

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
Haemophilus influenza type B	Epiglottitis Meningitis	Droplet	✓			✓ Until patient has completed 24 hours of antibiotics	For AGP * only	✓	Until 24 hours into the course of corrective antibiotic therapy.
Hepatitis, viral. Types A&E	Hepatitis	Contact		✓				✓	Hep A - For duration of hospital stay Hep E - For duration of hospital illness. Specifically for nappy wearing infants/incontinent adults.
Herpes Simplex (HSV1 and 2)	Oral herpes, genital herpes, neonatal disseminated herpes	Contact		✓		✓			Can infect oral mucosa (HSV1) or genital tract (HSV 2). Primary and recurrent infections can occur and duration of precautions will vary but usually until lesions or cold sores disappear. Risk to exposed infants delivered vaginally or by C-section and if mother has active infection and membranes have been ruptured for more than 4-6 hours

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
Herpes Zoster (Varicella zoster virus)	Shingles – skin	Contact	√ If lesions cannot be covered						<ul style="list-style-type: none"> http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/Shingles/ <p>Infectious until vesicles are dry usually 5-7 days this may be extended for immunocompromised individuals. Disseminated disease may require an increase in the duration of the precautions. Susceptible health/social care workers should not give direct care if immune care givers are available. Immune caregivers no additional precautions, non immune may require surgical mask in disseminated disease</p>
	Shingles – respiratory tract	Droplet/airborne	√			√	For AGP * only		
Influenza virus	Influenza	Droplet		√		√	For AGP * only	√	<p>DH UK Pandemic Influenza Preparedness Strategy 2015</p> <p>5 days except in immunocompromised persons. Use of vaccine or antiviral drugs may be considered. Avoid placing infected individuals with immunocompromised patients.</p>

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
Measles virus	Measles	Droplet / airborne			✓	✓	For AGP * only	✓	<p>PHE Measles guidance</p> <ul style="list-style-type: none"> http://www.hpa.org.uk/webc/HPAwebFile/HPAwebC/1274088429847 <p>Precautions remain until 4 days after onset of rash. For immunocompromised individuals this is increased for the duration of illness.</p> <p>Susceptible health/social care workers should not enter the room if immune care givers are available.</p> <p>Exposed susceptible care givers may require post exposure vaccine</p>
Multi drug resistant organisms	Various clinical syndromes, dependent upon organism and patient profile	Contact/Air borne/		✓					<p>PHE Carbapenemase Resistant Enterococci guidance</p> <ul style="list-style-type: none"> http://www.hpa.org.uk/webc/HPAwebFile/HPAwebC/1317140378646
Mumps virus	Mumps	Droplet		✓		✓	For AGP * only	✓	<p>Until approximately 9 days following appearance of symptoms in hospital. Some evidence that this can be reduced to 5 days in community settings for previously healthy individuals.</p> <p>Non immune HCW should not provide direct care</p>

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
Mycobacterium tuberculosis	Pulmonary/ laryngeal TB	Airborne			✓ Until patient has received 14 days effective treatment OR If patient has MDR or XDR TB		For AGP* only. Until patient has received 14 days effective treatment OR If patient has MDR or XDR TB	✓	NICE TB Guidelines 2011 – need new hyperlink <ul style="list-style-type: none"> http://www.nice.org.uk/nicemedia/live/13422/53642/53642.pdf Discontinue precautions only when patient is on effective therapy, condition is improving and has 3 negative sputum smears for acid fast bacilli (AFB) collected on 3 consecutive days. MDR TB cases always seek guidance when suspected/ confirmed cases further information (link to guidance) There are some exceptions to precaution requirements and local Infection Control/Health Protection/ TB teams must be consulted
	Extra pulmonary TB	Contact		✓		✓	For AGP* only	✓	
Mycoplasma	Pneumonia	Droplet		✓		✓	For AGP* only		Precautions remain for duration of hospital stay or until symptoms resolve. Patients can be infectious for up to 13 weeks.

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
<i>Neisseria meningitidis</i>	Meningitis Meningococcal septicaemia	Droplet	✓			✓ Until patient has completed 24 hours of antibiotics	For AGP * only	✓	Until 24 hours into the course of corrective antibiotic therapy. Post exposure chemoprophylaxis based on local risk assessments for exposed HCW as per Infection Control Team / Health Protection Team advice. Household contacts are given prophylactic antibiotics to eliminate carriage and prevent clinical illness,
Norovirus	Diarrhoea and vomiting	Contact/droplet	✓			Only if risk of splashing or spillage		✓	PHE guidance – need new hyperlink <ul style="list-style-type: none"> http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/Norovirus/Guidelines/
Parainfluenza virus	Respiratory tract infection	Droplet	✓			✓	For AGP * only		

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
Parvovirus B19	Erythema infectiosum/ 'slapped cheek' syndrome	Droplet		✓		In acute setting, up to 7 days from onset of symptoms	For AGP * only		A common childhood infection lasting 2- 3 days followed by the rash on the cheeks. In adults can be associated with arthralgia Non-infectious when the rash appears. If the patient has a chronic disease or is immunocompromised maintain precautions for the duration of illness or whilst patient is hospitalised. Advice should be sought from an Infection Control/Disease/ Public Health Physician or Consultant Microbiologist when this disease is identified or suspected in pregnancy
Respiratory syncytial virus (RSV)	Respiratory tract infection	Droplet		✓		✓	For AGP * only		DH guidance – greenbook hyperlink <ul style="list-style-type: none"> f PHE guidance <ul style="list-style-type: none"> http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/RespiratorySyncytialVirus/Guidelines/ Duration of symptoms (whilst in acute care setting specifically) Particularly affects young children, infants and immunosuppressed patients. Highly transmissible in paediatrics

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
Rhinovirus	Respiratory tract infection	Droplet		✓		✓	For AGP* only		
Rotaviruses	Gastroenteritis	Droplet/contact		✓		Only if risk of splashing or spillage			
Rubella virus	Rubella 'German measles'	Droplet		✓		*	For AGP* only	✓	Until 7 days after onset of rash. Susceptible HCW should not provide direct clinical care
Staphylococcus aureus	Scalded skin syndrome	Contact			If lesions cannot be covered			✓	
Meticillin resistant Staphylococcus	Infection	Contact		✓				✓	MRSA Screening - new hyperlinl <ul style="list-style-type: none"> http://wales.gov.uk/topics/health/cmo/publications/cmo/2013/mrsascreening/?lang=en Guidelines <ul style="list-style-type: none"> http://www.his.org.uk//resource_library/mrsa.cfm?cit_id=435&FAArea1=customWidgets.content_view_1&usecache=false
	Colonisation**								

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
<i>Streptococcus pyogenes</i> (Group A Strep)	Respiratory	Droplet	✓			✓ Until patient has completed 24 hours of antibiotics	For AGP * only		
	Wound, bacteraemia, meningitis, other metastatic infection	Contact	✓					✓	<p>PHE guidance</p> <ul style="list-style-type: none"> http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/StreptococcalInfections/Guidelines/ <p>GAS guidance</p> <p><i>Guidelines for prevention and control of group A streptococcal infection in acute healthcare and maternity settings in the UK 2016</i></p>

HYWEL DDA UNIVERSITY HEALTH BOARD

Pathogen	Clinical condition	Precaution category	Optimal Placement while patient is considered infectious			Respiratory precaution while patient is considered infectious		Notifiable	Useful links/additional comments
			Single	Single ambulance transfer	Single room with negative pressure	Surgical facemask	FFP3		
<i>Streptococcus pneumoniae</i>	Respiratory	Droplet	✓			✓ Until patient has completed 24 hours of antibiotics	For AGP * only	✓	PHE guidance <ul style="list-style-type: none"> http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/Pneumococcal/GuidelinesPneumococcal/ - need new hyperlink
	Wound, bacteraemia, meningitis, other metastatic infection	Contact	✓					✓	
Varicella virus	Chickenpox	Droplet/airborne			✓	✓	For AGP * only	✓	Viral rash in Pregnancy <ul style="list-style-type: none"> http://www.hpa.org.uk/webc/HPAwebFile/HPAwebC/1294740918985 <p>Until all lesions are dry and crusted. In immunocompromised individuals with varicella pneumonia prolonged precautions may be required. Susceptible health/social care workers (e.g. those who are pregnant or immunocompromised) should not enter the room if immune care givers are available.</p>

*Aerosol generating procedures

HYWEL DDA UNIVERSITY HEALTH BOARD

20. APPENDIX 3 – SUMMARY OF PRECAUTIONS TO MINIMISE THE SPREAD OF INFECTIONS BY THE DROPLET ROUTE



**VISITORS; PLEASE REPORT
TO NURSE IN CHARGE
BEFORE ENTERING THIS
ROOM**

DROPLET PRECAUTIONS FOR ISOLATED PATIENT

ALL HEALTHCARE WORKERS:



Hands

Decontaminate hands before entering this room.




Personal Protective Equipment

Wear disposable apron, fluid repellant surgical mask, eye/facial protection and gloves before entering this room.



Door Keep door closed

Risk assessed ☐
Door required to remain open.
Initials: Date:



Before leaving

Decontaminate equipment prior to removal from room.
Discard gloves, apron, eye/facial protection and fluid repellant surgical mask in healthcare waste bin.
Decontaminate hands.

HYWEL DDA UNIVERSITY HEALTH BOARD





21. APPENDIX 4 - SUMMARY OF PRECAUTIONS TO MINIMISE THE SPREAD OF INFECTIONS BY CONTACT



**VISITORS; PLEASE REPORT
TO NURSE IN CHARGE
BEFORE ENTERING THIS
ROOM**

CONTACT PRECAUTIONS FOR ISOLATED PATIENT

ALL HEALTHCARE WORKERS:

	Hands	Decontaminate hands before entering this room.
	Personal Protective Equipment	Wear disposable apron and gloves before entering this room.
	Door Keep door closed	Risk assessed <input type="checkbox"/> Door required to remain open. Initials: <input type="text"/> Date: <input type="text"/>
	Before leaving	Decontaminate equipment prior to removal from room. Discard gloves and apron in healthcare waste bin. Decontaminate hands.

HYWEL DDA UNIVERSITY HEALTH BOARD

22. APPENDIX 4 - SUMMARY OF PRECAUTIONS TO MINIMISE THE SPREAD OF INFECTIONS BY THE AIRBORNE ROUTE



**VISITORS; PLEASE REPORT
TO NURSE IN CHARGE
BEFORE ENTERING THIS
ROOM**

AIRBORNE PRECAUTIONS FOR ISOLATED PATIENT

ALL HEALTHCARE WORKERS:

	Hands	Decontaminate hands before entering this room.
	Personal Protective Equipment	Wear disposable apron/gown, FFP3 respirator, eye/face protection and gloves before entering this room.
	Door	Keep door closed.
	Before leaving	Decontaminate equipment prior to removal from room. Discard gloves, apron/gown and eye/face protection in healthcare waste bin. Decontaminate hands.
	After leaving	Remove respirator and discard in healthcare waste bin. Decontaminate hands.