Reference:	FOI.1503.19
Subject:	Urology departments' MRI use for prostate cancer diagnosis
Date of Request:	29 August 2019

## Requested

- 1. Do you routinely conduct prostate MRI (bpMRI/mpMRI) scans before first prostate biopsy as part of the initial diagnostic process? (please tick all that apply):
  - a. Yes, using T2-weighted, diffusion-weighted (multi-b ADC and high/long and dynamic contrast enhanced (DCE) sequences
  - b. Yes, using T2-weighted, diffusion-weighted (multi-b ADC and high/long b) sequences but not DCE
  - c. No but we refer to another provider (please provide details)
  - d. No (please provide details)
- 2. If yes, to 1a: What percentage of men with suspected prostate cancer receive mpMRI before biopsy as part of the initial diagnostic process?
- 3. If yes, to 1b: What percentage of men with suspected prostate cancer receive bpMRI before biopsy as part of the initial diagnostic process?
- 4. If you are based in Wales, we acknowledge that Health Boards have been developing prebiopsy mpMRI implementation plans in recent months to bring their practice into line with NICE guideline [NG131], Prostate cancer: diagnosis and management.

Please provide details of when you expect the following to be implemented for all eligible people:

- multiparametric MRI offered as the first-line investigation for people with suspected clinically localised prostate cancer; and
- People whose Likert score is 3 or more to be offered multiparametric MRI-influenced prostate biopsy.
- 5. What are your eligibility criteria/exclusion criteria for prostate MRI? (please tick all that apply):
  - a. Age (please provide details)
  - b. Symptoms (please provide details)
  - c. Life expectancy (please provide details)
  - d. Contra-indications (please provide details)
  - e. Other (please provide details)
- 6. Are you using results from the prostate MRI before biopsy to rule some men out of biopsy as part of the initial diagnostic process? (yes/no)
- 7. Do you biopsy all PI-RADS or LIKERT 3 scores?
  - a. Yes
  - b. No
  - c. Dependent on other patient histology
- 8. What threshold do you mostly use for ruling men out of biopsy?
  - a. PI-RADs 3 and above
  - b. LIKERT 3 and above

- c. PI-RADs 4 and above
- d. LIKERT 4 and above
- e. Varies depending on age (Please provide detail)
- f. Varies depending on other factors (Please provide detail)
- 9. What percentage of men do you estimate are ruled out of biopsy?
- 10. Have there been any changes to your prostate MRI capacity in the last year? (please choose all that apply):
  - a. An additional or new MRI scanner
  - b. Increased MRI scanner slots for prostate
  - c. Agreement to use Dynamic Contrast Enhancement
  - d. No longer using Dynamic Contrast Enhancement
  - e. A scanner/magnet upgrade
  - f. other (free text)
- 11. Has the number of radiologists at your trust/health board who report prostate MRI scans changed in the last year?
  - a. Increased
  - b. Decreased
  - c. Stayed the same
- 12. How many radiologists at your trust/health board report at least 250 prostate MRI scans per year?
- 13. Which of the following processes do you follow to manage men ruled out of an immediate biopsy, but with a raised PSA?
  - a. NICE Guidelines: prostate cancer diagnosis and management (NG131)
  - b. A local protocol (please provide details)
  - c. Other (please provide details)

## Response:

- 1. Hywel Dda University Health Board (UHB) confirms answer b: Using T2-weighted, diffusion-weighted (multi-b ADC and high/long b) sequences but not DCE.
- 2. Not applicable.
- 3. The percentage of men with suspected prostate cancer that receive bpMRI before biopsy is approximately 65% for all suspected prostate cancer referrals.
- 4. The UHB has implemented the pathway. All Pre-Biopsy MRI is discussed with specialist Uro-Radiologists, a biopsy is offered where indicated (including targeted biopsy).
- 5. The UHB's eligibility criteria/exclusion criteria for prostate MRI is as follows:
  - a. Patients over the age of 80 are not offered routine Pre-Biopsy MRI.
  - b. Symptoms are not a factor in requesting MRI, if UTI is ruled out.
  - c. Life expectancy 10 years or more.
  - d. Relative contraindications are, pacemaker, claustrophobia (all general contraindications for MRI apply).

- 6. The UHB confirms results from the prostate MRI before biopsy are used, to rule some men out of biopsy as part of the initial diagnostic process.
- 7. The UHB confirms answer c: Dependent on other patient factors, such as comorbidity.
- 8. The UHB confirms answer f: Dependent on other patient factors, such as comorbidity, PSA density and history of previous biopsy.
- 9. The audit for this year has not been completed at present therefore the UHB does not currently hold this information.
- 10. The UHB confirms answer a: A new replacement MRI scanner is due to be installed in February 2020, which will provide additional capacity.
- 11. The UHB confirms answer c: No changes to the number of radiologists in the UHB who report prostate MRI scans.
- 12. The UHB does not hold this information.
- 13. The UHB confirms answer a: NICE Guidelines: prostate cancer diagnosis and management (NG131) are followed to manage men ruled out of an immediate biopsy, but with a raised PSA.